

SHARP PITT INEQUALITY AND LOGARITHMIC UNCERTAINTY PRINCIPLE FOR DUNKL TRANSFORM IN L^2

D. V. GORBACHEV, V. I. IVANOV, AND S. YU. TIKHONOV

ABSTRACT. We prove sharp Pitt's inequality for the Dunkl transform in $L^2(\mathbb{R}^d)$ with the corresponding weights. As an application, we obtain the logarithmic uncertainty principle for the Dunkl transform.

1. INTRODUCTION

Let $\Gamma(t)$ be the gamma function, \mathbb{R}^d be the real space of d dimensions, equipped with a scalar product $\langle x, y \rangle$ and a norm $|x| = \sqrt{\langle x, x \rangle}$. Denote by $\mathcal{S}(\mathbb{R}^d)$ the Schwartz space on \mathbb{R}^d and by $L^2(\mathbb{R}^d)$ the Hilbert space of complex-valued functions endowed with a norm $\|f\|_2 = (\int_{\mathbb{R}^d} |f(x)|^2 dx)^{1/2}$. The Fourier transform is defined by

$$\widehat{f}(y) = (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(x) e^{-i\langle x, y \rangle} dx.$$

W. Beckner [2] proved the Pitt inequality for the Fourier transform

$$(1.1) \quad \| |y|^{-\beta} \widehat{f}(y) \|_2 \leq C(\beta) \| |x|^\beta f(x) \|_2, \quad f \in \mathcal{S}(\mathbb{R}^d), \quad 0 < \beta < d/2,$$

with sharp constant

$$C(\beta) = 2^{-\beta} \frac{\Gamma(\frac{1}{2}(\frac{d}{2} - \beta))}{\Gamma(\frac{1}{2}(\frac{d}{2} + \beta))}.$$

Noting that $\|(-\Delta)^{\beta/2} f(x)\|_2 = \| |x|^\beta \widehat{f}(x) \|_2$, Pitt's inequality can be viewed as a Hardy–Rellich inequality $\| |x|^{-\beta} f(x) \|_2 \leq C(\beta) \|(-\Delta)^{\beta/2} f(x)\|_2$; see the papers by D. Yafaev [19] and S. Eilertsen [7] for alternative proofs and extensions of (1.1).

For $\beta = 0$, (1.1) is the Plancherel theorem. If $\beta > 0$ there is no extremiser in inequality (1.1) and its sharpness can be obtained on the set of radial functions.

The proof of (1.1) in [2] is based on an equivalent integral realization as a Stein–Weiss fractional integral on \mathbb{R}^d . D. Yafaev in [19] used the following decomposition (see [17, Chapt. IV])

$$(1.2) \quad L^2(\mathbb{R}^d) = \sum_{n=0}^{\infty} \oplus \mathcal{R}_n^d,$$

Date: May 13, 2015.

1991 *Mathematics Subject Classification.* 42B10, 33C45, 33C52.

Key words and phrases. Dunkl transform, Pitt inequality, sharp constants, uncertainty principle.

The first and the second authors were partially supported by RFFI N 13-01-00045, Ministry of education and science of Russian Federation (N 5414GZ, N 1.1333.2014K), and Dmitry Zimin's Dynasty Foundation. The third author was partially supported by MTM 2011-27637, 2014 SGR 289, and RFFI 13-01-00043.

where \mathcal{R}_0^d is the space of radial function, and $\mathcal{R}_n^d = \mathcal{R}_0^d \otimes \mathcal{H}_n^d$ is the space of functions in \mathbb{R}^d that are products of radial functions and spherical harmonics of degree n . Thanks to this decomposition it is enough to study inequality (1.1) on the subsets of \mathcal{R}_n^d which are invariant under the Fourier transform.

In this paper, following [19] and using similar decomposition of the space $L^2(\mathbb{R}^d)$ with the Dunkl weight, we prove sharp Pitt's inequality for the Dunkl transform.

Let $R \subset \mathbb{R}^d$ be a root system, R_+ be the positive subsystem of R , and $k: R \rightarrow \mathbb{R}_+$ be a multiplicity function with the property that k is G -invariant. Here $G(R) \subset O(d)$ is a finite reflection group generated by reflections $\{\sigma_a: a \in R\}$, where σ_a is a reflection with respect to a hyperplane $\langle a, x \rangle = 0$.

Throughout this paper we let

$$v_k(x) = \prod_{a \in R_+} |\langle a, x \rangle|^{2k(a)}$$

denote the Dunkl weight. Moreover, $d\mu_k(x) = c_k v_k(x) dx$, where

$$c_k^{-1} = \int_{\mathbb{R}^d} e^{-|x|^2/2} v_k(x) dx$$

is the Macdonald–Mehta–Selberg integral. Let $L^2(\mathbb{R}^d, d\mu_k)$ be the Hilbert space of complex-valued functions endowed with a norm $\|f\|_{2, d\mu_k} = \left(\int_{\mathbb{R}^d} |f(x)|^2 d\mu_k(x) \right)^{1/2}$.

Introduced by C. F. Dunkl, a family of differential-difference operators (Dunkl's operators) associated with G and k are given by

$$D_j f(x) = \frac{\partial f(x)}{\partial x_j} + \sum_{a \in R_+} k(a) \langle a, e_j \rangle \frac{f(x) - f(\sigma_a x)}{\langle a, x \rangle}, \quad j = 1, \dots, d,$$

The Dunkl kernel $e_k(x, y) = E_k(x, iy)$ is the unique solution of the joint eigenvalue problem for the corresponding Dunkl operators:

$$D_j f(x) = iy_j f(x), \quad j = 1, \dots, d, \quad f(0) = 1.$$

Let us define the Dunkl transforms as follows

$$F_k(f)(y) = \int_{\mathbb{R}^d} f(x) \overline{e_k(x, y)} d\mu_k(x), \quad F_k^{-1}(f)(x) = F_k(f)(-x),$$

where $F_k(f)$ and $F_k^{-1}(f)$ are the direct and inverse transforms correspondingly (see, e.g., [12]). For $k \equiv 0$ we have $F_0(f) = \widehat{f}$.

Our goal is to study Pitt's inequality for the Dunkl transform

$$(1.3) \quad \| |y|^{-\beta} F_k(f)(y) \|_{2, d\mu_k} \leq C(\beta, k) \| |x|^\beta f(x) \|_{2, d\mu_k}, \quad f \in \mathcal{S}(\mathbb{R}^d),$$

with the sharp constant $C(\beta, k)$.

Let us first recall some known results on Pitt's inequality for the Hankel transform. Let $\lambda \geq -1/2$. Denote by $J_\lambda(t)$ the Bessel function of degree λ and by $j_\lambda(t) = 2^\lambda \Gamma(\lambda + 1) t^{-\lambda} J_\lambda(t)$ the normalized Bessel function. Setting

$$b_\lambda = \left(\int_0^\infty e^{-t^2/2} t^{2\lambda+1} dt \right)^{-1} = \frac{1}{2^\lambda \Gamma(\lambda + 1)}$$

and $d\nu_\lambda(r) = b_\lambda r^{2\lambda+1} dr$, we define $\|f\|_{2, d\nu_\lambda} = \left(\int_{\mathbb{R}_+} |f(r)|^2 d\nu_\lambda(r) \right)^{1/2}$.

The Hankel transform is defined by

$$H_\lambda(f)(\rho) = \int_{\mathbb{R}_+} f(r) j_\lambda(\rho r) d\nu_\lambda(r).$$

Note that $H_\lambda^{-1} = H_\lambda$ (see [3, 5, 8]). Pitt's inequality for the Hankel transform is written as

$$(1.4) \quad \|\rho^{-\beta} H_\lambda(f)(\rho)\|_{2, d\nu_\lambda} \leq c(\beta, \lambda) \|r^\beta f(r)\|_{2, d\nu_\lambda}, \quad f \in \mathcal{S}(\mathbb{R}_+),$$

where $c(\beta, \lambda)$ is the sharp constant in (1.4) and $\mathcal{S}(\mathbb{R}_+)$ is the Schwartz space on \mathbb{R}_+ . Note that if $f \in \mathcal{R}_0^d$, a study of the Hankel transform is of special interest since the Fourier transform of a radial function can be written as the Hankel transform.

L. De Carli [3] proved that $c(\beta, \lambda)$ is finite only if $0 \leq \beta < \lambda + 1$. In [3, 4] it was also studied the sharp constant in (L^p, L^q) Pitt's inequality for the Hankel transform of type (1.4) in the case of $1 \leq p \leq q \leq \infty$.

For $\lambda = d/2 - 1$, $d \in \mathbb{N}$, the constant $c(\beta, \lambda)$ was calculated by D. Yafaev [19], and in the general case by S. Omri [11]. The proof of Pitt's inequality in [11] is rather technical and uses the Stein–Weiss type estimate for the so-called B-Riesz potential operator. Following [19], we give a direct and simple proof of inequality (1.4) in Section 2.

Let $|k| = \sum_{a \in R_+} k(a)$ and $\lambda_k = d/2 - 1 + |k|$. For a radial function $f(r)$, $r = |x|$, Pitt's inequality for the Dunkl transform (1.3) corresponds to Pitt's inequality for the Hankel transform (1.4) with $\lambda = \lambda_k$. Therefore the condition

$$(1.5) \quad 0 \leq \beta < \lambda_k + 1$$

is necessary for $C(\beta, k) < \infty$. Our goal is to show that in fact $C(\beta, k) = c(\beta, \lambda_k)$ if condition (1.5) holds. This will be proved in Section 3. Moreover, in Section 4, we use Pitt's inequality for the Dunkl transform to obtain the logarithmic uncertainty principle (see [2, 11, 15]). It is worth mentioning that different uncertainty principles for the Dunkl transform have been recently studied in, e.g., [9, 10, 13, 14].

Note that for the one-dimensional Dunkl weight

$$v_\lambda(t) = |t|^{2\lambda+1}, \quad d\mu_\lambda(t) = \frac{v_\lambda(t) dt}{2^{\lambda+1} \Gamma(\lambda+1)}, \quad \lambda \geq -1/2,$$

and the corresponding Dunkl transform

$$F_\lambda(f)(s) = \int_{\mathbb{R}} f(t) \overline{e_\lambda(st)} |t|^{2\lambda+1} d\mu_\lambda(t), \quad e_\lambda(t) = j_\lambda(t) - i j'_\lambda(t),$$

F. Soltani [15] proved Pitt's inequality that can be equivalently written as

$$(1.6) \quad \| |s|^{-\beta} F_\lambda(f)(s) \|_{2, d\mu_\lambda} \leq \max \{c(\beta, \lambda), c(\beta, \lambda+1)\} \| |t|^\beta f(t) \|_{2, d\mu_\lambda}$$

for $f \in \mathcal{S}(\mathbb{R})$ and $0 \leq \beta < \lambda + 1$. Since $c(\beta, \lambda) \geq c(\beta, \lambda+1)$ (see [19]), then in fact (1.6) holds with the constant $c(\beta, \lambda)$ and therefore, we have in this case $C(\beta, k) = c(\beta, \lambda_k)$.

Finally, we remark that Pitt's inequality in L^2 for the multi-dimensional Dunkl transform has been recently established in [16] in the case of $\lambda_k - 1/2 < \beta < \lambda_k + 1$. The obtained constant is not sharp.

2. PITT'S INEQUALITY FOR HANKEL TRANSFORM

Theorem 2.1. *Let $\lambda \geq -1/2$ and $0 \leq \beta < \lambda + 1$, then for $f \in \mathcal{S}(\mathbb{R}_+)$ inequality (1.4) holds with the sharp constant*

$$c(\beta, \lambda) = 2^{-\beta} \frac{\Gamma(\frac{1}{2}(\lambda + 1 - \beta))}{\Gamma(\frac{1}{2}(\lambda + 1 + \beta))}.$$

Proof. For $\beta = 0$ we have $c(\beta, \lambda) = 1$ and (1.4) becomes Plancherel's identity

$$\|H_\lambda(f)(\rho)\|_{2, d\nu_\lambda} = \|f(r)\|_{2, d\nu_\lambda}.$$

Let $\beta > 0$. Setting $g(r) = f(r)r^{\beta+\lambda+1/2}$ in (1.4), we arrive at

$$\left(\int_0^\infty \left| \int_0^\infty g(r) J_\lambda(\rho r) (\rho r)^{1/2-\beta} dr \right|^2 d\rho \right)^{1/2} \leq c(\beta, \lambda) \left(\int_0^\infty |g(r)|^2 dr \right)^{1/2}.$$

Hence, $c(\beta, \lambda)$ is the norm of the integral operator

$$A_\lambda g(\rho) = \int_0^\infty a_\lambda(\rho r) g(r) dr : L^2(\mathbb{R}_+) \rightarrow L^2(\mathbb{R}_+),$$

where $a_\lambda(t) = J_\lambda(t)t^{1/2-\beta}$.

By [1, Sect. 7.7], the Mellin transform of the function $a_\lambda(\cdot)$ for $0 < \beta < \lambda + 1$ and $\eta \in \mathbb{R}$ is given by

$$\begin{aligned} Ma_\lambda(\eta) &= \int_0^\infty a(t)t^{-1/2-i\eta} dt = \int_0^\infty J_\lambda(t)t^{-\beta-i\eta} dt \\ &= 2^{-\beta-i\eta} \frac{\Gamma(\frac{1}{2}(\lambda+1-\beta-i\eta))}{\Gamma(\frac{1}{2}(\lambda+1+\beta+i\eta))}. \end{aligned}$$

Basic properties of the gamma function imply that $Ma_\lambda(\eta)$ is continuous and

$$Ma_\lambda(0) = 2^{-\beta} \frac{\Gamma(\frac{1}{2}(\lambda+1-\beta))}{\Gamma(\frac{1}{2}(\lambda+1+\beta))}, \quad |Ma_\lambda(\eta)| \sim 2^{-\beta} |\eta|^{-\beta} \rightarrow 0, \quad \eta \rightarrow \infty.$$

In [19, 20] it was proved that

$$\|A_\lambda\| = \max_{\mathbb{R}} |Ma_\lambda(\eta)| = Ma_\lambda(0)$$

and that the norm is not attained; that is to say, there is no function g such that $\|A_\lambda\| = \|A_\lambda g\|_{L^2(\mathbb{R}_+)}$ with $\|g\|_{L^2(\mathbb{R}_+)} = 1$. \square

Corollary 2.2. *If $\lambda_k = d/2 - 1 + |k|$ and $0 \leq \beta < \lambda_k + 1$, then for $f \in \mathcal{S}(\mathbb{R}^d) \cap \mathcal{R}_0^d$ Pitt's inequality for the Dunkl transform (1.3) holds with sharp constant $c(\beta, \lambda_k)$.*

3. PITT'S INEQUALITY FOR DUNKL TRANSFORM

Let \mathbb{S}^{d-1} be the unit sphere in \mathbb{R}^d , $x' \in \mathbb{S}^{d-1}$, and dx' be the Lebesgue measure on the sphere. Set $a_k^{-1} = \int_{\mathbb{S}^{d-1}} v_k(x') dx'$, $d\omega_k(x') = a_k v_k(x') dx'$, and $\|f\|_{2, d\omega_k} = \left(\int_{\mathbb{S}^{d-1}} |f(x')|^2 d\omega_k(x') \right)^{1/2}$. We have

$$c_k^{-1} = \int_{\mathbb{R}^d} e^{-|x|^2/2} v_k(x) dx = \int_0^\infty e^{-r^2/2} r^{d-1+2|k|} dr \int_{\mathbb{S}^{d-1}} v_k(x') dx' = b_{\lambda_k}^{-1} a_k^{-1}.$$

Let us denote by $\mathcal{H}_n^d(v_k)$ the subspace of k -spherical harmonics of degree $n \in \mathbb{Z}_+$ in $L^2(\mathbb{S}^{d-1}, d\omega_k)$ (see [6, Chap. 5]). Let $\Delta_k = \sum_{j=1}^d D_j^2$ be the Dunkl Laplacian and \mathcal{P}_n^d be the space of homogeneous polynomials of degree n in \mathbb{R}^d . Then $\mathcal{H}_n^d(v_k)$ is the restriction of $\ker \Delta_k \cap \mathcal{P}_n^d$ to the sphere \mathbb{S}^{d-1} .

If l_n is the dimension of $\mathcal{H}_n^d(v_k)$, we denote by $\{Y_n^j : j = 1, \dots, l_n\}$ the real-valued orthonormal basis $\mathcal{H}_n^d(v_k)$ in $L^2(\mathbb{S}^{d-1}, d\omega_k)$. A union of these bases forms

an orthonormal basis in $L^2(\mathbb{S}^{d-1}, d\omega_k)$ consisting of k -spherical harmonics, i.e., we have

$$(3.2) \quad L^2(\mathbb{S}^{d-1}, d\omega_k) = \sum_{n=0}^{\infty} \oplus \mathcal{H}_n^d(v_k).$$

Using (3.2) and the following Funk-Hecke formula for k -spherical harmonic $Y \in \mathcal{H}_n^d(v_k)$ (see [18])

$$(3.3) \quad \int_{\mathbb{S}^{d-1}} Y(y') \overline{e_k(x, y')} d\omega_k(y') = \frac{(-i)^n \Gamma(\lambda_k + 1)}{2^n \Gamma(n + \lambda_k + 1)} Y(x') r^n j_{n+\lambda_k}(r), \quad x = rx' \in \mathbb{R}^d,$$

similarly to (1.2) we have the direct sum decomposition of $L^2(\mathbb{R}^d, d\mu_k)$:

$$(3.4) \quad L^2(\mathbb{R}^d, d\mu_k) = \sum_{n=0}^{\infty} \oplus \mathcal{R}_n^d(v_k), \quad \mathcal{R}_n^d(v_k) = \mathcal{R}_0^d \otimes \mathcal{H}_n^d(v_k),$$

and that the space $\mathcal{R}_n^d(v_k)$ is invariant under the Dunkl transform. An example of the orthogonal basis in $L^2(\mathbb{R}^d, d\mu_k)$ consisting of eigenfunctions of the Dunkl transform was constructed in [6].

The next result provides a sharp constant in the Pitt inequality for the Dunkl transform (1.3).

Theorem 3.1. *Let $\lambda_k = d/2 - 1 + |k|$ and $0 \leq \beta < \lambda_k + 1$, then for $f \in \mathcal{S}(\mathbb{R}^d)$ we have*

$$C(\beta, k) = 2^{-\beta} \frac{\Gamma(\frac{1}{2}(\lambda_k + 1 - \beta))}{\Gamma(\frac{1}{2}(\lambda_k + 1 + \beta))}.$$

Sharpness of this inequality can be seen by considering radial functions.

Proof. For $\beta = 0$ we have $C(\beta, k) = 1$ and Pitt's inequality (1.3) becomes Plancherel's identity

$$\|F_k(f)(y)\|_{2, d\mu_k} = \|f(x)\|_{2, d\mu_k}.$$

Let $0 < \beta < \lambda_k + 1$. If $f \in \mathcal{S}(\mathbb{R}^d)$, then

$$f_{nj}(r) = \int_{\mathbb{S}^{d-1}} f(rx') Y_n^j(x') d\omega_k(x') \in \mathcal{S}(\mathbb{R}_+),$$

$$f(rx') = \sum_{n=0}^{\infty} \sum_{j=1}^{l_n} f_{nj}(r) Y_n^j(x'),$$

$$\int_{\mathbb{S}^{d-1}} |f(rx')|^2 d\omega_k(x') = \sum_{n=0}^{\infty} \sum_{j=1}^{l_n} |f_{nj}(r)|^2.$$

Using spherical coordinates, decomposition of $L^2(\mathbb{R}^d, d\mu_k)$ (3.4), formulas (3.1) and (3.3), and the property $e_k(tx, y) = e_k(ty, x)$, we get that

$$(3.5) \quad \begin{aligned} \int_{\mathbb{R}^d} |x|^{2\beta} |f(x)|^2 d\mu_k(x) &= b_{\lambda_k} \int_0^{\infty} r^{2\beta+d-1+2|k|} \int_{\mathbb{S}^{d-1}} |f(rx')|^2 d\omega_k(x') dr \\ &= b_{\lambda_k} \int_0^{\infty} r^{2\beta+d-1+2|k|} \sum_{n=0}^{\infty} \sum_{j=1}^{l_n} |f_{nj}(r)|^2 dr \\ &= \sum_{n=0}^{\infty} \sum_{j=1}^{l_n} \int_0^{\infty} |f_{nj}(r)|^2 r^{2\beta} d\nu_{\lambda_k}(r), \end{aligned}$$

$$\begin{aligned}
F_k(f)(y) &= \int_{\mathbb{R}^d} f(x) \overline{e_k(x, y)} d\mu_k(x) = b_{\lambda_k} \int_0^\infty r^{d-1+2|k|} \int_{\mathbb{S}^{d-1}} f(rx') d\omega_k(x') dr \\
&= \sum_{n=0}^\infty \sum_{j=1}^{l_n} b_{\lambda_k} \int_0^\infty f_{nj}(r) r^{2\lambda_k+1} \int_{\mathbb{S}^{d-1}} Y_n^j(x') \overline{e_k(rx', \rho y')} d\omega_k(x') dr \\
&= \sum_{n=0}^\infty \sum_{j=1}^{l_n} \frac{(-i)^n \Gamma(\lambda_k + 1)}{2^n \Gamma(n + \lambda_k + 1)} Y_n^j(y') \int_0^\infty f_{nj}(r) j_{n+\lambda_k}(\rho r) (\rho r)^n d\nu_{\lambda_k}(r),
\end{aligned}$$

and

$$\begin{aligned}
(3.6) \quad \int_{\mathbb{R}^d} |y|^{-2\beta} |F_k(f)(y)|^2 d\mu_k(y) &= \sum_{n=0}^\infty \sum_{j=1}^{l_n} \frac{\Gamma^2(\lambda_k + 1)}{2^{2n} \Gamma^2(n + \lambda_k + 1)} \\
&\quad \times \int_0^\infty \left| \int_0^\infty f_{nj}(r) j_{n+\lambda_k}(\rho r) (\rho r)^n d\nu_{\lambda_k}(r) \right|^2 \rho^{-2\beta} d\nu_{\lambda_k}(\rho).
\end{aligned}$$

Suppose that $g \in \mathcal{S}(\mathbb{R}_+)$, $n \in \mathbb{Z}_+$, and $0 < \beta < \lambda_k + 1 + n$. Let us show that

$$\begin{aligned}
(3.7) \quad \left\| \rho^{-\beta} \int_0^\infty g(r) j_{n+\lambda_k}(\rho r) (\rho r)^n d\nu_{\lambda_k}(r) \right\|_{2, d\nu_{\lambda_k}} \\
\leq \frac{2^n \Gamma(n + \lambda_k + 1) c(\beta, \lambda_k + n)}{\Gamma(\lambda_k + 1)} \|r^\beta g(r)\|_{2, d\nu_{\lambda_k}},
\end{aligned}$$

where $c(\beta, \lambda_k + n)$ is given in Theorem 2.1 with $\lambda = \lambda_k + n$. Setting in (3.7) $g(r) = u(r)r^n$, we rewrite it as follows:

$$(3.8) \quad \left\| \rho^{-\beta} \int_0^\infty u(r) j_{n+\lambda_k}(\rho r) d\nu_{\lambda_k+n}(r) \right\|_{2, d\nu_{\lambda_k+n}} \leq c(\beta, \lambda_k + n) \|r^\beta u(r)\|_{2, d\nu_{\lambda_k+n}},$$

which is (1.4) with $\lambda = \lambda_k + n$.

Since $c(\beta, \lambda_k + n)$ is decreasing with n (see [19]), then using (3.5), (3.6), and (3.7), we arrive at

$$\begin{aligned}
\int_{\mathbb{R}^d} |y|^{-2\beta} |F_k(f)(y)|^2 d\mu_k(y) &\leq \sum_{n=0}^\infty \sum_{j=1}^{l_n} c^2(\beta, \lambda_k + n) \int_0^\infty |f_{nj}(r)|^2 r^{2\beta} d\nu_{\lambda_k}(r) \\
&\leq c^2(\beta, \lambda_k) \sum_{n=0}^\infty \sum_{j=1}^{l_n} \int_0^\infty |f_{nj}(r)|^2 r^{2\beta} d\nu_{\lambda_k}(r) \\
&= c^2(\beta, \lambda_k) \int_{\mathbb{R}^d} |x|^{2\beta} |f(x)|^2 d\mu_k(x).
\end{aligned}$$

□

In the proof of Theorem 3.1 we obtained the following result (see (3.8)).

Corollary 3.2. *Let $n \in \mathbb{N}$, $\lambda_k = d/2 - 1 + |k|$, and $0 \leq \beta < \lambda_k + 1 + n$, then for $f \in \mathcal{S}(\mathbb{R}^d) \cap \mathcal{R}_n^d(v_k)$ we have Pitt's inequality for the Dunkl transform (1.3) with sharp constant $c(\beta, \lambda_k + n)$.*

For the Fourier transform Corollary 3.2 was established in [19].

4. LOGARITHMIC UNCERTAINTY PRINCIPLE FOR DUNKL TRANSFORM

W. Beckner in [2] proved the logarithmic uncertainty principle for the Fourier transform using Pitt's inequality (1.1): if $f \in \mathcal{S}(\mathbb{R}^d)$, then

$$\int_{\mathbb{R}^d} \ln(|x|)|f(x)|^2 dx + \int_{\mathbb{R}^d} \ln(|y|)|\widehat{f}(y)|^2 dy \geq \left(\psi\left(\frac{d}{4}\right) + \ln 2 \right) \int_{\mathbb{R}^d} |f(x)|^2 dx,$$

where $\psi(t) = d \ln \Gamma(t)/dt$ is the ψ -function.

For the Hankel transform the logarithmic uncertainty principle reads as follows (see [11]): if $f \in \mathcal{S}(\mathbb{R}_+)$ and $\lambda \geq -1/2$, then

$$(4.1) \quad \int_{\mathbb{R}_+} \ln(t)|f(t)|^2 t^{2\lambda+1} dt + \int_{\mathbb{R}_+} \ln(s)|H_\lambda(f)(s)|^2 s^{2\lambda+1} ds \\ \geq \left(\psi\left(\frac{\lambda+1}{2}\right) + \ln 2 \right) \int_{\mathbb{R}_+} |f(t)|^2 t^{2\lambda+1} dt.$$

For the one-dimensional Dunkl transform of functions $f \in \mathcal{S}(\mathbb{R})$, F. Soltani [15] has recently proved that

$$\int_{\mathbb{R}} \ln(|t|)|f(t)|^2 |t|^{2\lambda+1} dt + \int_{\mathbb{R}} \ln(|s|)|F_\lambda(f)(s)|^2 |s|^{2\lambda+1} ds \\ \geq \left(\min \left\{ \psi\left(\frac{\lambda+1}{2}\right), \psi\left(\frac{\lambda+2}{2}\right) \right\} + \ln 2 \right) \int_{\mathbb{R}} |f(t)|^2 |t|^{2\lambda+1} dt.$$

Since ψ is increasing the latter can be written as

$$(4.2) \quad \int_{\mathbb{R}} \ln(|t|)|f(t)|^2 |t|^{2\lambda+1} dt + \int_{\mathbb{R}} \ln(|s|)|F_\lambda(f)(s)|^2 |s|^{2\lambda+1} ds \\ \geq \left(\psi\left(\frac{\lambda+1}{2}\right) + \ln 2 \right) \int_{\mathbb{R}} |f(t)|^2 |t|^{2\lambda+1} dt,$$

which is the logarithmic uncertainty principle for the one-dimensional Dunkl transform.

Using Pitt's inequality (1.3) we obtain the logarithmic uncertainty principle the multi-dimensional Dunkl transform.

Theorem 4.1. *Let $\lambda_k = d/2 - 1 + |k|$ and $f \in \mathcal{S}(\mathbb{R}^d)$. We have*

$$(4.3) \quad \int_{\mathbb{R}^d} \ln(|x|)|f(x)|^2 d\mu_k(x) + \int_{\mathbb{R}^d} \ln(|y|)|F_k(f)(y)|^2 d\mu_k(y) \\ \geq \left(\psi\left(\frac{\lambda_k+1}{2}\right) + \ln 2 \right) \int_{\mathbb{R}^d} |f(x)|^2 d\mu_k(x).$$

Proof. We write the Pitt inequality (1.3) in the following form

$$\int_{\mathbb{R}^d} |y|^{-\beta} |F_k(f)(y)|^2 d\mu_k(y) \leq c^2(\beta/2, \lambda_k) \int_{\mathbb{R}^d} |x|^\beta |f(x)|^2 d\mu_k(x), \quad 0 \leq \beta < 2(\lambda_k+1).$$

For $\beta \in (-2(\lambda_k+1), 2(\lambda_k+1))$ define the function

$$\varphi(\beta) = \int_{\mathbb{R}^d} |y|^{-\beta} |F_k(f)(y)|^2 d\mu_k(y) - c^2(\beta/2, \lambda_k) \int_{\mathbb{R}^d} |x|^\beta |f(x)|^2 d\mu_k(x).$$

Since $|\beta| < 2(\lambda_k + 1)$ and $f, F_k(f) \in \mathcal{S}(\mathbb{R}^d)$, then

$$\int_{|x| \leq 1} |\ln(|x|)| |x|^\beta v_k(x) dx = \int_0^1 |\ln(r)| r^{\beta+2\lambda_k+1} dr \int_{\mathbb{S}^{d-1}} v_k(x') dx' < \infty,$$

which gives

$$|y|^{-\beta} \ln(|y|) |F_k(f)(y)|^2 v_k(y) \in L^1(\mathbb{R}^d) \quad \text{and} \quad \ln(|x|) |x|^\beta |f(x)|^2 v_k(x) \in L^1(\mathbb{R}^d).$$

Therefore,

$$(4.4) \quad \begin{aligned} \varphi'(\beta) = & - \int_{\mathbb{R}^d} |y|^{-\beta} \ln(|y|) |F_k(f)(y)|^2 d\mu_k(y) \\ & - c^2(\beta/2, \lambda_k) \int_{\mathbb{R}^d} |x|^\beta \ln(|x|) |f(x)|^2 d\mu_k(x) \\ & - \frac{dc^2(\beta/2, \lambda_k)}{d\beta} \int_{\mathbb{R}^d} |x|^\beta |f(x)|^2 d\mu_k(x). \end{aligned}$$

Pitt's inequality and Plancherel's theorem imply that $\varphi(\beta) \leq 0$ for $\beta > 0$ and $\varphi(0) = 0$ correspondingly, hence

$$\varphi'(0_+) = \lim_{\beta \rightarrow 0_+} \frac{\varphi(\beta) - \varphi(0)}{\beta} \leq 0.$$

Noting that

$$(4.5) \quad - \left. \frac{dc^2(\beta/2, \lambda_k)}{d\beta} \right|_{\beta=0} = \psi\left(\frac{\lambda_k + 1}{2}\right) + \ln 2,$$

we conclude that proof of (4.3) combining (4.4) and (4.5). \square

For the radial functions inequality (4.3) with $\lambda_k = \lambda$ implies inequality (4.1).

REFERENCES

- [1] H. Bateman and A. Erdélyi, Higher Transcendental Functions. Vol. 2, New York MacGraw-Hill, 1953.
- [2] W. Beckner, Pitt's inequality and uncertainty principle, *J. Proc. Amer. Math. Soc.*, **123** (1995), 1897–1905.
- [3] L. De Carli, On the L^p – L^q norm of the Hankel transform and related operators, *J. Math. Anal. Appl.*, **348** (1) (2008), 366–382.
- [4] L. De Carli, D. Gorbachev, and S. Tikhonov, Pitt and Boas inequalities for Fourier and Hankel transforms, *J. Math. Anal. Appl.*, **408** (2) (2013), 762–774.
- [5] L. Colzani, A. Crespi, L. Travaglini, and M. Vignati, Equiconvergence theorems for Fourier–Bessel expansions with applications to the harmonic analysis of radial functions in euclidean and noneuclidean spaces, *Trans. Amer. Math. Soc.*, **338** (1) (1975), 43–55.
- [6] C. F. Dunkl and Y. Xu, Orthogonal Polynomials of Several Variables. Cambridge Univ. Press. 2001.
- [7] S. Eilertsen, On weighted fractional integral inequalities, *J. Funct. Anal.*, **185** (1) (2001), 342–366.
- [8] A. Erdelyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, Tables of Integral Transforms. Vol. 2, McGraw-Hill Book Company, 1954.
- [9] S. Ghobber and P. Jaming, Uncertainty principles for integral operators, *Studia Math.*, **220** (2014), 197–220.
- [10] T. Kawazoe and H. Mejjaoli, Uncertainty principles for the Dunkl transform, *Hiroshima Math. J.*, **40** (2010), 241–268.
- [11] S. Omri, Logarithmic uncertainty principle for the Hankel transform, *Int. Trans. Spec. Funct.*, **22** (2011), 655–670.

- [12] M. Rösler, Dunkl operators. Theory and applications, in Orthogonal Polynomials and Special Functions, Lecture Notes in Math. Springer-Verlag, pp. 93–135, 2002.
- [13] M. Rösler, An uncertainty principle for the Dunkl transform, *Bull. Austral. Math. Soc.*, **59** (1999), 353–360.
- [14] N. Shimeno, A note on the uncertainty principle for the Dunkl transform, *J. Math. Sci. Univ. Tokyo.*, **8** (2001), 33–42.
- [15] F. Soltani, Pitt’s inequality and logarithmic uncertainty principle for the Dunkl transform on \mathbb{R} , *Acta Math. Hungar.*, **143** (2) (2014), 480–490.
- [16] F. Soltani, Pitt’s inequalities for the Dunkl transform on \mathbb{R}^d , *Int. Trans. Spec. Funct.*, **25** (9) (2014), 686–696.
- [17] E. Stein and G. Weiss, Introduction to Fourier Analysis on Euclidean Spaces. Princeton Univ. press, 1971.
- [18] Y. Xu, Dunkl operators: Funk–Hecke formula for orthogonal polynomials on spheres and on balls, *Bull. London Math. Soc.*, **32** (2000), 447–457.
- [19] D. Yafaev, Sharp constants in the Hardy–Rellich inequalities, *J. Funct. Anal.*, **168** (1) (1999), 121–144.
- [20] D. R. Yafaev, The discrete spectrum in the singular Friedrichs model, in Advances in the Mathematical Sciences, 189, Providence, pp. 255–274, Amer. Math. Soc. 1999.

D. GORBACHEV, TULA STATE UNIVERSITY, DEPARTMENT OF APPLIED MATHEMATICS AND COMPUTER SCIENCE, 300012 TULA, RUSSIA

E-mail address: `dvgmail@mail.ru`

V. IVANOV, TULA STATE UNIVERSITY, DEPARTMENT OF APPLIED MATHEMATICS AND COMPUTER SCIENCE, 300012 TULA, RUSSIA

E-mail address: `ivaleryi@mail.ru`

S. TIKHONOV, ICREA, CENTRE DE RECERCA MATEMÀTICA, AND UAB, CAMPUS DE BELLATERRA, EDIFICI C 08193 BELLATERRA (BARCELONA), SPAIN.

E-mail address: `stikhonov@crm.cat`